


considered. Molecular oxygen scavenges the hydrogen atoms, forming OOH radicals (eq 7) and lowering the recombination of OH[•] and H[•] radicals (eq 8).



Consequently more hydroxyl radicals are available if they escape from the bubble (high frequency) in the medium. At low frequency, in the long-lived bubble, OH[•] can be scavenged by reaction with the hydroperoxyl radical (eq 9); consequently H₂O₂ and I₃⁻ formation are more effective under argon.¹⁶



Changing the frequency of ultrasonic waves then produces modifications in the behavior of the species first formed in water sonolysis. This finding constitutes the first example of a clear frequency effect and should have important consequences for optimizing sonochemical oxidation yields.

Registry No. TMP, 2896-70-0; TMPone, 826-36-8; H₂O, 7732-18-5; O₂, 7782-44-7; HO[•], 3352-57-6; I⁻, 20461-54-5; I₃⁻, 14900-04-0; H₂O₂, 7722-84-1.

(16) (a) Sridaran, U. C.; Qlu, L. X.; Kaufman, F. *J. Phys. Chem.* **1982**, *86*, 4569. (b) Bielski, B. H. J.; Cabelli, D. E.; Arudi, R. L.; Ross, A. B. *J. Phys. Chem. Ref. Data* **1985**, *14*, 1041. (c) Atkinson, R. *Chem. Rev.* **1985**, *85*, 69. (d) Mozurkewich, M. *J. Phys. Chem.* **1986**, *90*, 2216.

Photoswitchable Binding of Substrates to Proteins: Photoregulated Binding of α -D-Mannopyranose to Concanavalin A Modified by a Thiophenefulgidye

Itamar Willner* and Shai Rubin

*Institute of Chemistry
The Hebrew University of Jerusalem
Jerusalem 91904, Israel*

Johann Wonner, Franz Effenberger, and Peter Bäuerle

*Institut für Organische Chemie und
Isotopenforschung der Universität Stuttgart
Pfaffenwaldring 55, 7000 Stuttgart 80, Germany*

Received January 3, 1992

Macromolecules exhibiting photoswitchable physical or chemical properties are extensively examined as information storage and signal amplification materials. Photoregulated "on-off" biomaterials provide a novel means to design targeted therapeutic agents activated and deactivated by external light signals. Various means to photoregulate biotransformations by light-switchable enzymes have been described and include the modification of the enzyme active site¹⁻⁴ and protein backbone^{5,6} by photochromic components and immobilization of enzymes in photochromic copolymers.⁷ Here we wish to report on the photoregulation of the binding properties of a protein by its chemical modification with photochromic units. We describe the photoswitchable binding of saccharides to concanavalin A modified by thiophenefulgidye.

Concanavalin A (Con A) is a globular lectin composed of four subunits (MW = 26K).⁸ Each subunit includes binding sites for

* Author to whom correspondence should be addressed.

(1) Bieth, J.; Vratisanos, S. M.; Wassermann, N. H.; Erlanger, B. F. *Proc. Natl. Acad. Sci. U.S.A.* **1969**, *64*, 1103.

(2) Wainberg, M. A.; Erlanger, B. F. *Biochemistry* **1971**, *10*, 3816.

(3) Turner, A. D.; Pizzo, S. V.; Rozakis, G. W.; Porter, N. A. *J. Am. Chem. Soc.* **1987**, *109*, 1274-1275.

(4) Porter, N. A.; Bruhuke, J. O. *Photochem. Photobiol.* **1990**, *51*, 37-43.

(5) Aizawa, M.; Namba, K.; Suzuki, S. *Arch. Biochem. Biophys.* **1977**, *182*, 305-310.

(6) Willner, I.; Rubin, S.; Riklin, A. *J. Am. Chem. Soc.* **1991**, *113*, 3321-3325.

(7) Willner, I.; Rubin, S.; Zor, T. *J. Am. Chem. Soc.* **1991**, *113*, 4013-4014.




Figure 1. 2-modified Con A: (a) 3-E obtained after irradiation of 3-C at $\lambda > 475$ nm; (b) 3-C obtained after irradiation of 3-E at 400 nm $> \lambda > 300$ nm.

Scheme I




Table I. Association Constants of 4 to 2-Modified Con A in Its Two Photochromic States (E and C) as a Function of Loading Degree by the Photochromic Material

loading degree ^a	dye configuration	$K_a \times 10^{-4} \text{ (M}^{-1}\text{)}^b$
0		2.2
6	E	1.64
6	C	2
9	E	0.78
9	C	1.21
12	E or C	0.64

^a Loading degree is defined as the number of photochromic components anchored to each protein backbone. The total number of lysine residues in Con A corresponds to 12, and thus loading degree corresponds to the number of substituted lysine residues. ^b $K_a = [(\text{Con A}) \cdot (4)] / [\text{Con A}][4]$. Association constants were determined by equilibrating Con A or 3-E or 3-C (30 μM) with 10 μM 4 in phosphate buffer solution (pH 7) of 0.0875 M, 0.1 mM CaCl₂, 0.1 mM MnCl₂, and 0.1 M NaCl at 20 °C. The protein was filtered off, and the concentration of free 4 was determined by HPLC (RP-18 column, eluent water/methanol, 50:50, at flow rate 1 mL/min, UV detection).

Mn²⁺ and Ca²⁺ and monosaccharide binding sites.^{9,10} The metal ions act cooperatively in the association of the monosaccharides to Con A. The affinity of Con A toward monosaccharides is sensitive to the steric configuration of the pyranose hydroxyl groups at positions C-3, C-4, and C-6 of the saccharide.¹¹ Two pyranoses, α -D-mannopyranose and α -D-glucopyranose, are recognized for binding by Con A, and the affinity of Con A toward α -D-mannopyranose is ca. 3.5-fold higher than toward α -D-glucopyranose.¹² Thus, small steric alterations of the protein backbone

(8) Edelman, G. M.; Cunningham, B. A.; Reeke, G. N.; Becker, J. W.; Waxdal, M. J.; Wang, J. L. *Proc. Natl. Acad. Sci. U.S.A.* **1972**, *69*, 2580-2584.

(9) Becker, J. W.; Reeke, G. N.; Wang, J. L.; Cunningham, B. A.; Edelman, G. M. *J. Biol. Chem.* **1975**, *250*, 1513-1524.

(10) Goldstein, I. J. *Carbohydrate-Protein Interaction*; American Chemical Society: Washington, D.C., 1979.

(11) Goldstein, I. J.; Hollerman, C. E.; Smith, E. E. *Biochemistry* **1965**, *4*, 876.

(12) Hassing, G. S.; Goldstein, I. J. *Eur. J. Biochem.* **1970**, *16*, 549-556.





Figure 2. Reversible photoregulated complexation of 2-modified Con A to 4. Points 2 and 4 (O) correspond to complexation states of the substrate when 2-modified Con A is in the *C* form. Points 1, 3, and 5 (●) correspond to decomplexation states when 2-modified Con A is in the *E* form.

are anticipated to affect its binding properties toward these pyranoses.


Fulgide dyes are well established photochromic materials.¹³ Thiophenfulgide (1)¹⁴ undergoes three photostimulated isomerization processes¹⁵ (eq 1) where UV irradiation of 1-*Z* results in 1-*E* that undergoes subsequent conrotatory electrocyclicization to 1-*C*. Reversible isomerization of 1-*C* to 1-*E* is photoinduced



by filtered light, $\lambda > 475$ nm.¹⁶ Con A (Sigma) was modified by thiophenfulgimide (2).¹⁷ The modification process was carried out by reacting 50 mg of Con A in 6 mL of an aqueous solution that contained 250 mg NaHCO_3 with 5–15 mg of 2 dissolved in 200 μL of THF at 0 °C for 24–48 h (Scheme 1). This modification leads to functionalization of Con A lysine residues, and the loading degree is controlled by the amount of added 2 and the reaction time. The resulting modified protein, 3, exhibits photochromic properties (Figure 1). Illumination of 3-*E*, 300 nm $< \lambda < 400$ nm, results in isomerization to 3-*C*. Further illumination of 3-*C*, $\lambda > 475$ nm, restores the original absorption spectrum of 3-*E*. The loading degrees of Con A by 2 are determined by following the absorbance of 3-*C* at $\lambda = 532$ nm ($\epsilon = 2200 \text{ M}^{-1} \text{ cm}^{-1}$) and determining the protein content or by comparing the fluorescence intensities resulting from the interaction of fluorescamine¹⁸ with 3-*E* and unmodified Con A. Both methods give excellent agreement ($\pm 5\%$).

Photoregulated association of 4-nitrophenyl α -D-mannopyranoside (4) to 3-*E* and 3-*C* has been followed by determination of the association constants of 4 to the photochromic protein.

Table I summarizes the association constants of 4 to Con A in its two photochromic states as a function of loading degree. It is evident that, as the loading degree of Con A increases, its



affinity for 4 declines and the association constant decreases in its value. It is also concluded that, up to a loading degree of 9, the affinity of 3-*C* is higher than that of 3-*E* toward binding of 4. At a loading degree of 12, where all lysine residues of Con A are modified by 2, a substantial decrease in the binding constant of 4 is observed, and no difference between states *E* and *C* is detected. The largest difference in binding constants of 3-*E* and 3-*C* to 4 is observed at a loading degree of 9. This difference in the binding constants allows us to reversibly photoswitch the association and dissociation of 4 to 2-modified Con A (Figure 2). We see that in the presence of 3-*E*, only 2.4×10^{-6} M of 4 is bound. Upon illumination, $\lambda = 300$ –400 nm, and isomerization to 3-*C*, enhanced binding of 4 is observed. Further illumination, $\lambda > 475$ nm, regenerates 3-*E* and the intermolecular complex is dissociated.

We thus conclude that chemical modification of Con A by thiophenfulgimide (2) allows the photoinduced binding of 4-nitrophenyl α -D-mannopyranoside to the protein. Photoswitchable association and dissociation of the protein–substrate assembly is induced by the two photochromic states of the protein. Further studies on photoregulation of the binding of substrates to other proteins using this approach are in progress in our laboratory.

Acknowledgment. This research was supported by the Basic Research Foundation, administered by the Israel Academy of Sciences and Humanities, and the Deutsche Forschungsgemeinschaft (SFB 329).

Registry No. Thiophenfulgide, 118408-54-1; α -D-mannopyranose, 7296-15-3.

Reaction of (6*R*)-6-F-EPSP with Recombinant *Escherichia coli* Chorismate Synthase Generates a Stable Flavin Mononucleotide Semiquinone Radical

Manoj N. Ramjee,[†] Shankar Balasubramanian,[†] Chris Abell,^{*‡} John R. Coggins,[§] Gareth M. Davies,^{||} Timothy R. Hawkes,[‡] David J. Lowe,[†] and Roger N. F. Thorneley^{*†}

*AFRC Institute of Plant Science Research
Nitrogen Fixation Laboratory, University of Sussex
Brighton, Sussex, BN1 9RQ, UK*

*Department of Chemistry, University Chemical
Laboratory, University of Cambridge*

Lensfield Road, Cambridge, CB2 1EW, UK

*Department of Biochemistry, University of Glasgow
Glasgow, G12 8QQ, UK*

*ICI Pharmaceuticals, Mereside, Alderley Park
Macclesfield, Cheshire, SK10 4TG, UK*

*ICI Agrochemicals, Jealotts Hill Research Station
Bracknell, Berkshire, RG12 6EY, UK*

Received December 16, 1991

Chorismate synthase (EC 4.6.1.4), the seventh enzyme in the shikimate pathway,¹ catalyzes the conversion of 5-enolpyruvylshikimate 3-phosphate (1, EPSP) to chorismate 2.² The reaction involves the removal of the C-6 *pro-R* hydrogen and loss of

(13) *Photochromism: Molecules and Systems*; Dürr, H.; Bouas-Larent, H. Eds.; Elsevier: Amsterdam, 1990.

(14) Glaze, A. P.; Harris, S. A.; Heller, H. G.; Johncock, W.; Oliver, S. N.; Strydom, P. J.; Whittall, J. *J. Chem. Soc., Perkin Trans. 1* **1985**, 957.

(15) Ulrich, K.; Port, H.; Wolf, H. C.; Wonner, J.; Effenberger, F.; Ilge, H. D. *Chem. Phys.* **1991**, *154*, 311.

(16) Ulrich, K.; Port, H.; Bäuerle, P. *Chem. Phys. Lett.* **1984**, *155*, 437.

(17) The synthesis of 2 will be published elsewhere.

(18) Udenfriend, S.; Stein, S.; Bohlen, P.; Dairman, W.; Leimgruber, W.; Weigle, M. *Science* **1972**, *178*, 871.

* Authors to whom correspondence should be addressed.

[†] University of Sussex.

[‡] University of Cambridge.

[§] ICI Pharmaceuticals.

^{||} ICI Agrochemicals.

(1) Bentley, R. *Crit. Rev. Biochem. Mol. Biol.* **1990**, *25* (5), 307–384.

(2) Welch, G. R.; Cole, K. W.; Gaertner, F. H. *Arch. Biochem. Biophys.* **1974**, *165*, 505–518.