

(Iminomethylidene)phosphines (RP=C=NR)

Curt Wentrup,*^{1a} Horst Briehl,^{1a} Gerd Becker,^{1a,b}
Gudrun Uhl,^{1a,b} Hans-Jürgen Wessely,^{1a,b}
André Maquestiau,^{1c} and Robert Flammang^{1c}

Department of Chemistry, University of Marburg
D-3550 Marburg, West Germany
Institute for Inorganic Chemistry
University of Stuttgart
D-7000 Stuttgart 80, West Germany
Laboratory of Organic Chemistry
State University of Mons
B-7000 Mons, Belgium
Received August 15, 1983

There has been much recent interest in compounds incorporating P=C^{2,3} and P=P⁴ double bonds. Furthermore, the stable (2,2-dimethylpropylidene)phosphine, (CH₃)₃CC≡P, featuring a PC triple bond has been prepared.⁵ However, previous attempts to isolate cumulenes possessing a two-coordinate phosphorus atom, RP=C=X, have failed,^{2,3,6} except in a single case, the preparation of the stable and sterically protected (CH₃)₃CP=C=NC(CH₃)₃.⁷ We now wish to report a general method of preparation of unstable (iminomethylidene)phosphines (monophosphorus analogues of carbodiimides), their direct observation by IR and mass spectrometry, and their initial chemical reactions.

The required starting materials, (iminomethyl)phosphines **1**, exist in equilibrium with carbamoylphosphines [(CH₃)₃SiP(R)-CON(Ph)Si(CH₃)₃] and smaller amounts of alkylidenephosphines RP=C(OSi(CH₃)₃)N(Ph)Si(CH₃)₃.⁸ In analogy with the reaction used in the preparation of (CH₃)₃CC≡P,⁵ the compounds **1** could be expected to eliminate hexamethyldisiloxane and thus give rise to the desired (iminomethylidene)phosphines, RP=C=NPh (Scheme I).







Figure 1. IR spectrum (-196 °C) of the product of pyrolysis of **3a** at 480 °C.

Scheme I



In the event, the treatment of **1** with a catalytic amount of solid NaOH led only to the isolation of the dimers **3**.⁹ **3b** and **3e** were also obtained by spontaneous decomposition of the adducts **2b** and

(9) (a) Satisfactory elemental analyses (C, H, N, P) were obtained for these compounds. (b) X-ray crystallographic determinations of **3c** and **3d** confirm the structure assignments. The two exocyclic phosphorus ligands are trans oriented. Full crystallographic data will be published: Becker, G.; Härer, J.; Riffel, H.; Uhl, G.; Wessely, H.-J., in preparation.

(1) (a) University of Marburg. (b) Present address: University of Stuttgart. (c) University of Mons.

(2) Appel, R.; Knoll, F.; Ruppert, I. *Angew. Chem.* **1981**, *93*, 771; *Angew. Chem., Int. Ed. Engl.* **1981**, *20*, 731.

(3) Becker, G.; Becker, W.; Mundt, O. *Phosphorus Sulfur* **1983**, *14*, 267.

(4) Yoshifuji, M.; Shima, I.; Inamoto, N.; Hirotsu, K.; Higuchi, T. *J. Am. Chem. Soc.* **1981**, *103*, 4587. Cowley, A. H.; Kilduff, J. E.; Newman, T. H.; Pakulski, M. *Ibid.* **1982**, *104*, 5820. Bertrand, G.; Couret, C.; Escudie, J.; Majid, S.; Majoral, J.-P. *Tetrahedron Lett.* **1982**, *23*, 3567. Couret, C.; Escudie, J.; Satgé, J. *Ibid.* **1982**, *23*, 4941. Centinkaya, B.; Hudson, A.; Lappert, M. F.; Goldwhite, H. J. *J. Chem. Soc., Chem. Commun.* **1982**, 609. Centinkaya, B.; Hitchcock, P. B.; Lappert, M. F.; Thorne, A. J.; Goldwhite, H. *Ibid.* **1982**, 691. Niecke, E.; Rüger, R.; Lysek, M.; Pohl, S.; Schoeller, W. *Angew. Chem.* **1983**, *95*, 495; *Angew. Chem., Int. Ed. Engl.* **1983**, *22*, 486; Cowley, A. H.; Kilduff, J. E.; Norman, N. C.; Pakulski, M.; Atwood, J. L.; Hunter, W. E. *J. Am. Chem. Soc.* **1983**, *105*, 4845.

(5) Becker, G.; Gresser, G.; Uhl, W. *Z. Naturforsch., B* **1981**, *36B*, 16.

(6) Appel, R.; Laubach, B. *Tetrahedron Lett.* **1980**, *21*, 2497.

(7) Kolodiaznyii, O. I. *Tetrahedron Lett.* **1982**, *23*, 4933; *J. Gen. Chem. USSR (Engl. Transl.)* **1982**, *52*, 2101.

(8) Itoh, K.; Fukui, M.; Ishi, Y. *J. Chem. Soc. C* **1969**, 2002. Becker, G.; Mundt, O. *Z. Anorg. Allg. Chem.* **1979**, *459*, 87. Issleib, K.; Schmidt, H.; Meyer, H. *J. Organomet. Chem.* **1980**, *192*, 33. Becker, G.; Wessely, H.-J., manuscript in preparation.




Figure 2. CID mass spectrum of m/z 211 from the pyrolysis of **3a** at 430 °C.

2e, the latter formed by the addition of phenyl isothiocyanate to the alkyl- or aryl(trimethylsilyl)phosphine.¹⁰

If the diphosphetanes **3** are, in fact, dimers of (iminomethylidene)phosphines, the monomers might be regenerated via a cycloreversion under suitable reaction conditions. This, indeed, is readily achieved by flash vacuum pyrolysis of **3**. Product formation was monitored by low-temperature IR spectroscopy, and a pyrolysis unit has also been attached to the ion source of a reversed-geometry Varian MAT 311A mass spectrometer¹¹ equipped with a collision chamber for obtaining CID (collisionally induced dissociation) spectra of initial pyrolysis products.

2,4-Bis(phenylimino)-1,3-diphenyl-1,3-diphosphetane (**3a**) (mp 139 °C) was vaporized at 139–145 °C (10^{-5} torr). When the pyrolysis temperature was increased to 400 °C, an IR band at 1853 cm^{-1} appeared in the pyrolyzate condensed at -196 °C. The maximum intensity of this band was observed at a pyrolysis temperature at 480 °C (Figure 1). On warming the pyrolyzate to -55 °C, the intensity of the 1853- cm^{-1} band started decreasing; at the same time, a band at 1560 cm^{-1} due to the starting material **3a** started increasing. This process was complete at -30 °C, and after warming to room temperature **3a** was recovered in better than 90% yield.¹²

These observations indicate that **3a** dissociates into two molecules of the (iminomethylidene)phosphine **4a** on gas-phase pyrolysis. The monomeric **4a** reverts to **3a** above -55 °C in the solid state. The strong band at 1853 cm^{-1} in the IR of **4a** is assigned to the asymmetric stretching vibration of the $\text{P}=\text{C}=\text{N}$ moiety. For comparison, carbodiimides ($\text{RN}=\text{C}=\text{NR}$) absorb near 2100 cm^{-1} .

The analogous pyrolysis of **3b** (mp 220 °C) at 480 °C (sublimation temperature 110 °C, increasing to 200 °C in 35 min) gave rise to a strong IR band at 1839 cm^{-1} ascribed to **4b**. This material started redimerizing to **3b** at -25 °C, a process that was complete at 0 °C. Comparable results were obtained with the precursors **3c–e**.

The IR assignments were corroborated by using the mass spectrometry reactor. With increasing pyrolysis temperature, the

M^+ peak due to **3a** decreased, while that of **4a** increased. The CID mass spectrum of **4a** obtained at 430 °C is shown in Figure 2. The base peak at m/z 183 may be ascribed to the dibenzophospholyl cation, $\text{C}_{12}\text{H}_8\text{P}^+$, which is typical of diphenylated phosphorus compounds.¹³ In the pyrolysis of **3b** the parent peak of the precursor (m/z 382) disappeared at 490 °C, while that of **4b** (m/z 191) reached maximum intensity. Here, the spectrum is dominated by a loss of isobutene to give m/z 135, formally corresponding to $\text{PhN}=\text{C}=\text{Ph}^+$ or $\text{PhNH}-\text{C}\equiv\text{P}^+$.

We are continuing the studies of (iminomethylidene)phosphines, in particular cycloaddition reactions and attempts to obtain other phosphorus-containing cumulenes, e.g., $\text{RP}=\text{C}=\text{O}$.

Registry No. **1a**, 24103-42-2; **1b**, 87218-80-2; **1c**, 87729-47-3; **1d**, 87729-48-4; **1e**, 87729-49-5; **2b**, 87729-50-8; **2e**, 87729-51-9; **3a**, 87729-52-0; **3b**, 87218-81-3; **3c**, 87729-53-1; **3d**, 87729-54-2; **3e**, 87729-55-3; **4a**, 87729-56-4; **4b**, 87218-77-7; **4c**, 87729-57-5; **4d**, 87729-58-6; **4e**, 87729-59-7.

(13) Budzikiewicz, H.; Djerassi, C.; Williams, D. H. "Mass Spectrometry of Organic Compounds"; Holden-Day: San Francisco, 1967; p 647.

(10) Becker, G.; Uhl, G., unpublished results.

(11) Maquestiau, A.; Van Haverbeke, Y.; De Meyer, C.; Flammang, R.; Perlaux, J. *Bull. Soc. Chim. Belg.* **1976**, *85*, 69.

(12) The band at 2120 cm^{-1} in Figure 1 is due to phenyl isocyanide, formed in a competing thermal fragmentation of **3a**. This material evaporates during warm-up and thus does not contaminate the final product. A yield of ca. 10% of phenyl isocyanide was obtained by distilling it into a cold trap and subsequently identifying it by comparison with an authentic sample.