

THE TRANSPOSITION OF LOCALLY COMPACT,  
CONNECTED TRANSLATION PLANES

Thomas Buchanan and Hermann Hähl

This note deals with the transposition of translation planes in the topological context. We show that a topological congruence  $C$  of the real vector space  $\mathbb{R}^{2n}$  has the property that every hyperplane of  $\mathbb{R}^{2n}$  contains a component of  $C$ . This makes it possible to define the transpose  $P^T$  of the topological translation plane  $P$  associated with  $C$ ; it is proved that the translation plane  $P^T$  is topological also. The relationship between collineation groups and the relationship between coordinatizing quasifields of  $P$  and  $P^T$  are also discussed.

§1.

Before we formulate our main results, we recall some facts and definitions.

The affine point set of a locally compact, connected, topological translation plane  $P$  can be identified topologically with a finite dimensional real vector space  $V$  (in its usual topology) in such a manner that the lines through the origin of  $V$  are linear subspaces ([3, p.12-13]). These lines form a congruence of  $V$ ; and the translation plane  $P$  is the translation plane associated with this congruence in the sense of André [1]. Any congruence of a real finite dimensional vector space  $V$  which can be obtained in this manner will be called a *topological congruence* of  $V$ . It is known that the only possible dimensions of  $V$  are 2, 4, 8 and 16 ([3, p.47 Satz 1]).

By  $V^*$  we denote the dual space of  $V$  consisting of

all linear forms  $V \rightarrow \mathbb{R}$ ; by the *polar* of a linear subspace  $C$  of  $V$  we mean as usual the annihilator  $C^0 = \{\varphi \in V^* / \varphi(C) = 0\}$  of  $C$  in  $V^*$ .

Using results of Breuning [3] we shall prove

**THEOREM.** *Let  $C$  be a topological congruence of a real finite dimensional vector space  $V$ . Then*

(a)  *$C$  has the property that any hyperplane of  $V$  contains a component of  $C$ , and*

(b) *the set  $C^0 = \{C^0 / C \in C\}$  of the polars of the components of  $C$ , which by (a) is a congruence of  $V^*$ , is a topological congruence.*

*Remark:* The topological hypothesis of our theorem cannot be omitted, since by [2, Teorema 2.2] there exist (necessarily nontopological) congruences of  $V$  which do not have the hyperplane property of part (a).

**COROLLARY.** *Let  $P$  be a locally compact, connected, topological translation plane, and let  $C$  be the associated topological congruence of the underlying real vector space  $V$ . Then the translation plane  $P^\tau$  associated with the corresponding polar congruence  $C^0$  is also a locally compact, connected, topological translation plane.*

The plane  $P^\tau$  is called the *transpose*<sup>1)</sup> of  $P$ .

*Proof of the Theorem.*

(a) Identify  $V$  with  $\mathbb{R}^{2n}$  and give  $C$  the finest topology such that the map

$$(1) \quad \mathbb{R}^{2n} \setminus \{0\} \rightarrow C: x \mapsto C_x,$$

which sends each nonzero vector  $x$  to the unique component  $C_x$  of  $C$  containing  $x$ , is continuous. By Satz 2 of [3, p.14]  $C$  is homeomorphic to the  $n$ -sphere  $S^n$ .

---

1) The authors are indebted to Dr. Rainer Löwen for calling their attention to this elegant description of the transpose using polars.

Moreover, if we consider  $C$  as a subset of the Grassmann manifold  $G_n(\mathbb{R}^{2n})$  of  $n$ -dimensional subspaces of  $\mathbb{R}^{2n}$ , then the topology of  $C$  is the topology induced from  $G_n(\mathbb{R}^{2n})$  ([3, p.30 Satz 1 and p.17]).

Let  $H$  be an arbitrary but fixed hyperplane in  $\mathbb{R}^{2n}$ . Arguing indirectly, we assume that  $H$  does not contain a component of  $C$ . Then obviously  $n \geq 2$ . Moreover, every component of  $C$  would intersect  $H$  in a linear subspace of dimension  $n-1$ . Thus

$$C|H = \{C \cap H \mid C \in C\}$$

would be a subset of the Grassmann manifold  $G_{n-1}(H)$ . By looking at the Grassmann manifolds as quotient spaces of Stiefel manifolds it is easy to see that the map

$$(2) \quad C \longrightarrow G_{n-1}(H): C \longmapsto C \cap H$$

would be continuous. Thus  $C|H$  as a subspace of  $G_{n-1}(H)$  would be homeomorphic to  $S^n$ .

Consider the canonical  $(n-1)$ -dimensional vector bundle  $\gamma^{n-1}(H)$  over  $G_{n-1}(H)$  as defined for example in Milnor-Stasheff [6, p.59 ff.]. By restricting  $\gamma^{n-1}(H)$  to the subspace  $C|H$  of  $G_{n-1}(H)$  and by removing the zero vector from each fiber, we would obtain a locally trivial fiber bundle over the  $n$ -sphere  $C|H$  with fiber  $\mathbb{R}^{n-1} \setminus \{0\}$ , total space

$$E = \{(C \cap H, x) \in (C|H) \times (H \setminus \{0\}) \mid x \in C\},$$

and bundle projection  $E \longrightarrow C|H$  given by the projection onto the first factor. The total space  $E$  would be homeomorphic to  $H \setminus \{0\} = \mathbb{R}^{2n-1} \setminus \{0\}$  by the map  $H \setminus \{0\} \longrightarrow E: x \longmapsto (C_x \cap H, x)$ , which can be expressed using maps (1) and (2) to check continuity. We now note an inconsistency when we examine the following part of the exact homotopy sequence of this fiber bundle:

$$\longrightarrow \pi_{n-1}(S^n) \xrightarrow{\partial} \pi_{n-2}(\mathbb{R}^{n-1} \setminus \{0\}) \longrightarrow \pi_{n-2}(\mathbb{R}^{2n-1} \setminus \{0\}) \longrightarrow ;$$

since  $n \geq 2$ , the homotopy groups (or sets)  $\pi_{n-1}(S^n)$  and  $\pi_{n-2}(\mathbb{R}^{2n-1} \setminus \{0\})$  are zero, whereas  $\pi_{n-2}(\mathbb{R}^{n-1} \setminus \{0\})$  is not zero. This contradiction proves part (a) of the theorem.

(b) By [6, p.57 Lemma 5.1] the map  $C \mapsto C^\circ$  from  $C$  into the Grassmann manifold  $G_n(V^*)$  is continuous, since after appropriate identification of the vector spaces  $V$  and  $V^*$ , the polar  $C^\circ$  of  $C$  is nothing more than the orthogonal complement  $C^\perp$ . Therefore  $C^\circ$  is homeomorphic to  $S^n$  too.

Thus to complete the proof of part (b) it suffices to apply the following lemma, which is implicit in [3].

*LEMMA. Let  $C$  be a congruence of  $\mathbb{R}^{2n}$  having linear subspaces as components. Then  $C$  is topological if and only if  $C$  is compact as a subset of  $G_n(\mathbb{R}^{2n})$ .*

The "only if" part of this lemma has already been used in part (a) of the proof; it follows immediately from [3, p.14 Satz 2, p.30 Satz 1 and p.17]. For the "if" part, we look at the canonical  $n$ -dimensional vector bundle  $\gamma^n(\mathbb{R}^{2n})$  restricted to  $C$  having the total space

$$E = \{(C,x) \in C \times \mathbb{R}^{2n} \mid x \in C\}$$

and the projection  $E \rightarrow C: (C,x) \mapsto C$ . This vector bundle admits (in the terminology of [3, p.22]) the effective Gauss map  $E \rightarrow \mathbb{R}^{2n}: (C,x) \mapsto x$ . We can then apply [3, p.27 Satz, p.28 Korollar and p.34 Satz 1].

## §2.

In this section we make a few remarks about the effects of transposition of a locally compact, connected translation plane as far as the collineation groups and coordinatizing quasifields are concerned.

Recall that a quasifield which coordinatizes a locally compact, connected translation plane  $P$  is a locally compact, connected, topological quasifield in the sense of Salzmann [7, §7]; conversely, any plane coordinatized by such a quasifield is a locally compact, connected translation plane. The kernel  $K$  of  $P$  is isomorphic as a topological field to  $\mathbb{R}$ ,  $\mathbb{C}$  or the quaternion field  $\mathbb{H}$  ([3, p.8 Satz 3]).

We first claim that the transpose  $P^t$  of  $P$  can be constructed equally well using  $K$  instead of  $R$  as the base field.

More generally let  $P$  be a (not necessarily topological) translation plane which is finite dimensional over its kernel  $K$ . By [1] the affine point set  $V$  of  $P$  may be considered as a one-sided vector space over  $K$  in such a manner that the components of the congruence  $C$  associated with  $P$  are  $K$ -linear subspaces. (In the case where  $P$  is locally compact and connected, the  $R$ -linear structure which we have been considering up to now is, of course, simply given by restriction to the closed subfield  $R$  of  $K$ .) Given any subfield  $K'$  of  $K$ , we can regard  $V$  as a  $K'$ -vector space and form its dual

$$V_{K'}^* = \text{Hom}_{K'}(V, K')$$

over  $K'$ . For  $C \in \mathcal{C}$  consider the  $K'$ -polar

$$C_{K'}^{\circ} = \{\psi \in V_{K'}^* \mid \psi(C) = 0\}$$

and the family

$$C_{K'}^{\circ} = \{C_{K'}^{\circ} \mid C \in \mathcal{C}\}.$$

**PROPOSITION 1.** *Suppose  $K'$  is a subfield of  $K$  and  $F$  a central subfield of  $K'$  such that  $V$  has finite dimension over all these fields. Then there exists an  $F$ -linear isomorphism  $V_{K'}^* \rightarrow V_F^*$  which maps  $C_{K'}^{\circ}$  onto  $C_F^{\circ}$  for every  $C \in \mathcal{C}$ . In particular, if one of the families  $C_F^{\circ}$  and  $C_{K'}^{\circ}$  is a congruence, the other is also.*

*Proof:* If  $\delta: K' \rightarrow F$  is any nonzero  $F$ -linear form, let

$$\Delta: V_{K'}^* \rightarrow V_F^*: \psi \mapsto \delta \circ \psi$$

be the induced map, which is  $F$ -linear, since  $F$  is central in  $K'$ . The map  $\Delta$  is injective, and  $\Delta(C_{K'}^{\circ}) \subseteq C_F^{\circ}$  for every  $C \in \mathcal{C}$ . Since the  $F$ -dimensions of  $V_F^*$  and  $V_{K'}^*$  are equal, and the  $F$ -dimensions of  $C_F^{\circ}$  and  $C_{K'}^{\circ}$  are equal,  $\Delta$  must be an isomorphism, and  $\Delta(C_{K'}^{\circ}) = C_F^{\circ}$ .

**COROLLARY.** *Let  $P$  be a locally compact, connected*

translation plane,  $K$  its kernel and  $C$  the congruence associated with  $P$ . Then  $C_K^{\circ}$  is a topological congruence, and the translation plane associated with  $C_K^{\circ}$  is topologically isomorphic to the transpose  $P^{\tau}$  described in the Corollary on page 2.

This follows from the Theorem on page 2 using Proposition 1 with  $F = \mathbb{R}$  and  $K' = K$ .

Maduram [5] has described the relationships between some collineation groups and between coordinatizing quasi-fields of finite translation planes and their transposes. His general results as well as his proofs carry over to locally compact, connected translation planes. We give topological analogues of three of his propositions, which are of particular interest in this context, together with indications of proofs using our notation.

PROPOSITION 2 (cf. [5, Prop. 3]). *Let  $P$  be a locally compact, connected translation plane, let  $\Gamma_0$  denote the isotropy subgroup of the affine collineation group of  $P$  with respect to the origin, and let  $\Sigma_0$  denote the subgroup of continuous collineations in  $\Gamma_0$ ; let  $\Gamma_0^*$  and  $\Sigma_0^*$  be the analogously defined groups of collineations of the transpose  $P^{\tau}$ . Then  $\Gamma_0$  and  $\Gamma_0^*$  are isomorphic--viewed as transformation groups of the congruences associated with  $P$  and  $P^{\tau}$  respectively--via an isomorphism which maps  $\Sigma_0$  onto  $\Sigma_0^*$ . The action of  $\Sigma_0^*$  on the translation group of  $P^{\tau}$  is the contragredient representation of the  $\mathbb{R}$ -linear action of  $\Sigma_0$  on the translation group of  $P$ .*

For the proof, recall that  $\Gamma_0$  consists of all  $K$ -semilinear automorphisms of  $V$  which respect the congruence  $C$  ([1, p.178 Satz 19]). For  $A \in \Gamma_0$  with  $\alpha \in \text{Aut } K$  its companion automorphism, define a  $K$ -semilinear transformation  $A^{\vee}$  of  $V_K^*$  by

$$(3) \quad A^{\vee}(\psi) = \alpha \circ \psi \circ A^{-1} \quad (\psi \in V_K^*).$$

A straightforward calculation shows that  $A^\vee$  respects the congruence  $C_K^\circ$ , and thus by the corollary on page 5 may be considered as an element of  $\Gamma_0^*$ . One easily checks that  $A \mapsto A^\vee: \Gamma_0 \rightarrow \Gamma_0^*$  is a group isomorphism; this and the bijection  $C \rightarrow C_K^\circ: C \mapsto C_K^\circ$  together constitute a transformation group isomorphism

$$(4) \quad (\Gamma_0, C) \xrightarrow{\cong} (\Gamma_0^*, C_K^\circ): (A, C) \mapsto (A^\vee, C_K^\circ).$$

Since the companion automorphism of  $A^\vee$  is again  $\alpha$ , the continuity of  $A$ ,  $\alpha$  and  $A^\vee$  are equivalent.

To compare the actions of  $\Sigma_0$  and  $\Sigma_0^*$ , take the  $\mathbb{R}$ -linear form  $\delta: K \rightarrow \mathbb{R}$  which associates to each  $z \in K$  its real part  $\delta(z) = \operatorname{Re} z$  ( $K$  is either  $\mathbb{R}$ ,  $\mathbb{C}$  or  $\mathbb{H}$ ), and identify  $V_K^*$  and  $V_{\mathbb{R}}^*$  by the induced isomorphism  $\Delta$  as defined in the proof of Proposition 1. We shall interpret formula (3) in  $V_{\mathbb{R}}^*$ : Making use of the fact that  $\delta \circ \alpha = \delta$  when  $\alpha$  is a continuous automorphism of  $K$ , we have

$$\Delta \circ A^\vee \circ \Delta^{-1}(\varphi) = \delta \circ \alpha \circ (\Delta^{-1}\varphi) \circ A^{-1} = \varphi \circ A^{-1} \quad (A \in \Sigma_0, \varphi \in V_{\mathbb{R}}^*),$$

which is the expression defining the contragredient representation.

PROPOSITION 3 (cf. [5, Prop. 4 and 5(3)]). *Let  $Q$  be a locally compact, connected quasifield (or a real finite dimensional division algebra), and let  $P$  be the translation plane coordinatized by  $Q$ . Then there is a locally compact, connected quasifield (respectively, a real division algebra)  $Q'$ , which coordinatizes the transpose  $P^T$  and thus has the same dimension as  $Q$ , such that the middle nucleus and the left nucleus of  $Q'$  are topologically isomorphic to the left nucleus and the middle nucleus of  $Q$  respectively.*

*Remark:* Note that Dembowski, André and Maduram consider quasifields with the distributive law  $(a + b)c = ac + bc$ , whereas the distributive law used by Breuning and Salzmann is  $c(a + b) = ca + cb$ . Proposition 3 is formulated ac-

ording to the latter convention; in particular, the left nucleus here plays the role of the right nucleus in [4, p. 134] and [5].

For the proof of the proposition, let  $C_1$  and  $C_2$  denote the first and second coordinate axes in the coordinatization of  $P$  over  $Q$ , and call their respective points at infinity  $p_1$  and  $p_2$ . Take  $Q'$  to be a locally compact, connected quasifield which coordinatizes  $P^\Gamma$  with respect to the  $K$ -polars  $(C_i)_K^O \in C_K^O$  ( $i = 1, 2$ ) as first and second coordinate axes. To simplify notation we write these axes as  $C_i^O$ ; denote their respective points at infinity by  $p_i^O$  ( $i = 1, 2$ ).

By straightforward calculation we see that the transformation group isomorphism  $(\Gamma_0, C) \cong (\Gamma_0^*, C_K^O)$  of (4) restricts to an isomorphism between the group  $\Gamma(p_i, C_j)$  of central collineations of  $P$  with center  $p_i$  and axis  $C_j$  and the group  $\Gamma^*(p_j^O, C_i^O)$  of central collineations of  $P^{\Gamma_j}$  with center  $p_j^O$  and axis  $C_i^O$  ( $i, j \in \{1, 2\}$ ). These groups of central collineations are  $K$ -linear ([1, p. 180 Satz 22]); hence the groups carry the obvious topology, and the isomorphisms above are clearly continuous. The proof proceeds by using the well-known relationships of these groups to substructures of the coordinatizing quasifields (cf. [4, p.134, 3.1.30 ff.]): The left-middle interchange in the assertion regarding the nuclei of  $Q$  and  $Q'$  follows from the isomorphisms  $\Gamma(p_1, C_2) \cong \Gamma^*(p_2^O, C_1^O)$  and  $\Gamma(p_2, C_1) \cong \Gamma^*(p_1^O, C_2^O)$  and the relationships of these strain groups to nuclei. Moreover,  $Q'$  is a semifield when  $Q$  is, since the shears group  $\Gamma(p_2, C_2)$  is transitive on  $C \setminus \{C_2\}$  if and only if the corresponding group  $\Gamma^*(p_2^O, C_2^O)$  is transitive on  $C_K^O \setminus \{C_2^O\}$ . Finally, a locally compact, connected semifield is a real finite dimensional (not necessarily associative) division algebra, because by the arguments of [3, p.8 Satz 3] it contains  $\mathbb{R}$  in its center.

The contribution of the second author to this paper is part of his work on a program sponsored by the *Deutsche Forschungsgemeinschaft*, and he would like to thank the *DFG*



for their support.

## REFERENCES

1. André, J.: Über nicht-desarguessche Ebenen mit transitiver Translationsgruppe. Math.Z. 60 (1954), 156-186.
2. Bernardi, M.: Esistenza di fibrazioni in uno spazio proiettivo infinito. Ist. Lombardo Acad.Sci.Lett.Rend. A 107 (1973), 528-542.
3. Breuning, P.: Translationsebenen und Vektorraumbündel. Mitt.Math.Sem. Giessen 86 (1970), 1-50.
4. Dembowski, P.: Finite Geometries. Berlin-Heidelberg-New York, Springer-Verlag 1968.
5. Maduram, D.M.: Transposed translation planes. Proc. Amer.Math.Soc. 53 (1975), 265-270.
6. Milnor, J., Stasheff, J.D.: Characteristic Classes. Princeton, Princeton University Press 1974.
7. Salzmann, H.: Topological planes. Advances in Math. 2 (1967), 1-60.

Thomas Buchanan, Hermann Hähl  
Mathematisches Institut  
der Universität Tübingen  
Auf der Morgenstelle 10  
D-7400 Tübingen 1  
Federal Republic of Germany

(Eingegangen am 3. Februar 1977)