

Photoelectron Spectra and Molecular Properties. XXX.¹ π Interactions in H_3Si - and H_3C -Substituted Acetylenes

Walter Ensslin,² Hans Bock,* and Gerd Becker

Contribution from the Institutes of Inorganic Chemistry, University of Frankfurt, Frankfurt-am-Main, Germany, and University of Karlsruhe, Karlsruhe, Germany. Received September 24, 1973

Abstract: The photoelectron (pe) spectra of mono- and disubstituted silyl- and methylacetylenes are assigned by comparison with the corresponding ionization potentials of acetylene, disilane, and ethane. The observed π splittings can be rationalized within a parameterized hyperconjugation model. Calculations using a modified CNDO/2 procedure with and without inclusion of 3d orbitals in the basis set demonstrate that the usual parameters tend to overemphasize $p_\pi d_\pi$ back-bonding. To rationalize available experimental data, 3d_{si} orbitals need not to be considered explicitly—neither for the different dipole moments of methyl- and silylacetylenes nor for the general pe spectroscopic pattern of silicon compounds, *i.e.*, high first ionization potential, reduced spin-orbit coupling, or band broadening.

The cylindrical π system of acetylene and its perturbation by substituents have been the subject of numerous investigations. In particular, the different properties of trimethylsilyl- and *tert*-butylacetylenes³ have been attributed to differences in inductive and conjugative effects of silyl and alkyl groups. In contrast, Pitt,⁴ on the basis of CNDO/2 calculations, proposed a hyperconjugative model neglecting $p_\pi d_\pi$ back-bonding to account for silicon substituent effects. Heilbronner, *et al.*,⁵ completely assigned the photoelectron spectra of trimethylsilylacetylene and its halogen derivatives and derived spectroscopically basis orbital parameters for the trimethylsilyl group. In the following we report the assignment and interpretation of the photoelectron spectra of silyl- and methylacetylenes $H_3XC\equiv CH$ and $H_3XC\equiv CXH_3$ ($X = C, Si$).

Pe Spectra and Hyperconjugation Model

The spectra of the four silyl- and methyl-substituted acetylenes are shown in Figure 1, and their vertical ionization potentials are compiled in Table I.

Based on Koopmans theorem⁶ the pe spectrum of acetylene⁷ has been interpreted in terms of the orbital sequence $(1\pi_u)^4 (3\sigma_g)^2 (2\sigma_u)^2$. The first band, corresponding to electron ejection from the doubly degenerate π_u orbital, consists of well-resolved peaks with $\Delta IP = 1830 \text{ cm}^{-1}$, the small reduction relative to the ground-state $C\equiv C$ stretching frequency $\nu_{C=C} = 1983 \text{ cm}^{-1}$ indicating only a modest change of the triple bond lengths in the radical cation.

In the pe spectrum of methylacetylene⁷ (Figure 1), the first pe band rises steeply and clearly displays the

acetylene π vibrational fine structure ($\nu_{C=C}$: ion, 1940 cm^{-1} ; ground state, 2142 cm^{-1}). In contrast, the π band of silylacetylene (C_{3v} : 2e) is shifted to higher ionization potential and considerably broadened and its vibrational spacing ($\nu_{C=C}$: ion, 1760 cm^{-1} ; ground state,⁸ 2192 cm^{-1}) is less pronounced. Such differences are frequently observed for alkyl-silyl exchange at π^9 or lone pair centers¹⁰ and are attributed to increasing π -type delocalization.⁶ At higher ionization potentials three pe bands are observed for methylacetylene and six for the silyl compound. Obviously, the ionization energies of silylacetylene except the first one are more or less lowered relative to those of methylacetylene (Figure 1 and Table I).

For a complete assignment we use a qualitative hyperconjugation model based on linear combinations of bond orbitals (LCBO) as specified in Figure 2. Applying second-order perturbation arguments to each set of symmetry equivalent basis orbitals yields a qualitative orbital sequence, in which one σ orbital, $4a_1$, is inserted between the π orbitals 2e and 1e, resulting from hyperconjugative mixing and characterized by a nodal plane in the threefold molecular axis. Within the LCBO MO model, the hyperconjugative interaction in methylacetylene can be parameterized identifying the energy of the basis orbital $e_{C=C}(\pi)$ with the first vertical ionization potential of acetylene $IP_{\pi^v} = 11.40 \text{ eV}$ and $e_{XH_3}(\pi)$ with the center of gravity of the ionization potentials of either the orbital pair e_g/e_u of H_3XXH_3 or the t_2 orbitals of XH_4 . For the parameterization of the silylacetylene hyperconjugation model, we start with disilane, the pe spectrum¹¹ of which shows two Jahn-Teller split bands, $1e_g$ and $1e_u$, at 11.88–12.13 and 12.73–13.08 eV. From the average value $e_{SiH_3}(\pi) = 12.45 \text{ eV}$ and the acetylene π ionization potential $e_{C=C}(\pi) = 11.40 \text{ eV}$, we obtain as the π center of

* Correspondence should be addressed to Institute of Inorganic Chemistry, 6 Frankfurt, Germany, Theodor-Stern-Kai 7.

(1) Part XXVIII: K. Wittel and H. Bock, *Chem. Ber.*, **107**, 317 (1974).

(2) W. Ensslin, Thesis, University of Frankfurt, 1974.

(3) H. Bock and H. Seidl, *J. Chem. Soc. B*, 1158 (1968), and literature quotations given therein.

(4) C. G. Pitt, *J. Organometal. Chem.*, **61**, 49 (1973), and literature quotations given therein.

(5) G. Bieri, F. Brogli, E. Heilbronner, and E. Kloster-Jensen, *J. Electron Spectrosc. Relat. Phenomena*, **1**, 67 (1972).

(6) *Cf.* for instance C. R. Brundle and M. B. Robin in "Determination of Organic Structures," Vol. III, F. C. Nachod and J. J. Zuckerman, Ed., Academic Press, New York, N. Y., 1971.

(7) D. W. Turner, C. Baker, A. D. Baker, and C. R. Brundle, "Molecular Photoelectron Spectroscopy," Wiley-Interscience, London, 1970.

(8) H. Siebert, "Anwendungen der Schwingungsspektroskopie in der anorganischen Chemie," Springer, Berlin, 1966, and literature quotations given therein.

(9) *Cf.* for instance P. Mollère, H. Bock, G. Becker, and G. Fritz, *J. Organometal. Chem.*, **46**, 89 (1972).

(10) *Cf.* for instance S. Craddock and E. A. V. Ebsworth, *Chem. Commun.*, 57 (1971); S. Craddock and R. A. Whiteford, *Trans. Faraday Soc.*, **67**, 3425 (1971); *J. Chem. Soc., Faraday Trans. 2*, **68**, 281 (1972); H. Bock, P. Mollère, G. Becker, and G. Fritz, *J. Organometal. Chem.*, **61**, 113, 127 (1973).

(11) H. Bock, W. Ensslin, F. Feher, and W. Freund, unpublished results.

Table I. Vertical Ionization Energies (eV) of Methyl- and Silylacetylenes, Acetylene, Ethylene, and Disilane

Compd	$\pi_{C=C}$	1	2	3	4	5	6
$H_3CC\equiv CH^6$	10.37		14.6	$\begin{pmatrix} 15.3 \\ 15.5 \end{pmatrix}_{sh}^a$	17.4		
$H_3SiC\equiv CH$	10.73		12.53	$\begin{pmatrix} 13.05 \\ 13.85 \end{pmatrix}_{J/T}^a$	17.4	18.9	20.4
$H_3CC\equiv CCH_3^{23}$	9.61		$(14.0)_{sh}$	14.5	$(15.0)_{sh}$	$(15.8)_{sh}$	20.63
$H_3SiC\equiv CSiH_3$	10.46		$(12.4)_{sh}$	12.9	14.51	18.2	19.0
$HC\equiv CH^7$	11.40		16.75	18.75			
$H_3CCH_3^7$		12.1	$\begin{pmatrix} 12.7 \\ 13.25 \end{pmatrix}_{J/T}^a$	$\begin{pmatrix} 15.5 \\ 15.7 \end{pmatrix}_{J/T}^a$	20.5		
$H_3SiSiH_3^2$		10.53	$\begin{pmatrix} 11.88 \\ 12.13 \end{pmatrix}_{J/T}^a$	$\begin{pmatrix} 12.73 \\ 13.08 \end{pmatrix}_{J/T}^a$	16.5		

^a J/T: π bands obviously split by the Jahn-Teller effect (see text).




Figure 1. Pe spectra of mono- and disubstituted methyl- and silylacetylenes.

gravity $\bar{e}(\pi) = 11.93$ eV. In the pe spectrum of silylacetylene (Figure 1) the corresponding mean, $\bar{e}(\pi) = [e_{SiH_3}(\pi) + e_{C=C}(\pi)]/2 = (13.45 + 10.73)/2 = 12.09$ eV, is shifted by +0.16 eV, probably because all unoccupied orbitals are neglected¹² within the LCBO MO model (see Figure 2). This value is used to correct the basis orbital energies $e_{SiH_3}(\pi) = 12.45 + 0.16 = 12.61$ eV and $e_{C=C}(\pi) = 11.40 + 0.16 = 11.56$ eV, henceforth called coulomb parameters $\alpha_{C=C}$ and α_{XH_3} . The LCBO interaction term, *i.e.*, the resonance integral $\beta_{\pi, H_3SiCCH}^{LCBO}$, is then calculated from the corresponding second-order determinant to be -1.25 eV. Analogously, with $e_{CH_3}(\pi) = 14.25$ eV (\bar{e} (ethane) = 14.3 eV and \bar{t}_2 (methane) = 14.2 eV¹³) one obtains

(12) Cf., *e.g.*, M. Beez, G. Bieri, H. Bock, and E. Heilbronner, *Helv. Chim. Acta*, **56**, 1028 (1973); H. Bock, J. Sauer, D. Seebach, G. Wagner, and K. Wittel, *Chem. Ber.*, in press.




Figure 2. LCBO MO model for monosubstituted acetylenes $H_3XC\equiv CH$.

for methylacetylene a hyperconjugation parameter $\beta_{\pi, H_3CCH}^{LCBO} = -2.07$ eV. As shown subsequently, these β_{π}^{LCBO} values allow calculation of the π ionization potentials of dimethyl- and disilylacetylenes as well.¹⁴ Therefore, the pe spectra of the monosubstituted acetylenes are tentatively assigned following the proposal in Figure 2. The second pe band corresponds to an ionization from the σ orbital $4a_1$ mainly representing the X-C bond, which is supported by the 2 eV reduction on methyl-silyl exchange or by the vibrational fine structures (*e.g.*, silyl deformation:³ ground state, 936 cm^{-1} ; ion, 890 cm^{-1}). The third band is due to electron ejection from the degenerate H_3X orbital $1e$ in accordance with the observed Jahn-Teller split and with the lack of vibrational fine structure. It is

(13) A. W. Potts and W. C. Price, *Proc. Roy. Soc., Ser. A*, **326**, 165 (1972).

(14) For restrictions, see F. Brogli, E. Heilbronner, V. Hornung, and E. Kloster-Jensen, *Helv. Chim. Acta*, in press.




Figure 3. Hyperconjugation model for methyl- and silylacetylenes, including center of gravity corrections (\downarrow) as well as pe ionization potentials of the $e(\pi)$ assigned bands. Observed Jahn-Teller splits are indicated by $\{ \}$ and assignments of overlapping bands according to parameterization by $()_{\text{calc}}$.

lowered in the silyl derivative by 2 eV. In contrast the fourth band stays almost constant representing more or less the acetylene C-H bond $3a_1$.

The parameter $\beta_{\pi}^{\text{LCBO}}$, calculated from $\text{H}_3\text{XC}\equiv\text{CH}$ ionization potentials, will now be utilized to assign the $\text{H}_3\text{XC}\equiv\text{CXH}_3$ pe spectra. To do so, our model has to be expanded for two e_{XH_3} basis orbitals, assuming free rotation in the $\text{H}_3\text{XC}\equiv\text{CXH}_3$ molecules and therefore the D_3 conformation being the most frequent one. From the linear combinations of the two e_{XH_3} group orbitals

$$\phi_{\text{XH}_3}^{\pm} = \frac{1}{\sqrt{2}}(\phi_1(e_{\text{XH}_3}) \pm \phi_2(e_{\text{XH}_3})) \quad (1)$$

only $\phi_{\text{XH}_3}^+$ will interact with the acetylene π orbital $\phi_{\text{C}\equiv\text{C}}$. Therefore only a split $\phi_{\text{XH}_3}^+/\phi_{\text{C}\equiv\text{C}}$ is to be expected, whereas $\phi_{\text{XH}_3}^-$ should remain unaffected. Starting with the coulomb parameters $\alpha_{\text{C}\equiv\text{C}} = -11.72$ eV (again corrected by +0.16 eV) and $\alpha_{\text{SiH}_3} = -12.77$ eV and with the interaction term $\beta_{\pi, \text{H}_3\text{SiC}\equiv\text{C}\text{SiH}_3}^{\text{LCBO}} = \sqrt{2}\beta_{\pi, \text{H}_3\text{SiCCH}}^{\text{LCBO}} = \sqrt{2}(-1.25) = -1.77$ eV, one calculates for disilylacetylene the π -orbital energies $\epsilon_{1,2} = -12.24 \pm 1.84$ eV. Of the resulting two eigenvalues, $\epsilon_1 = 10.4$ eV matches perfectly with the observed π ionization potential $IP_{\pi}^{\nu} = 10.46$ eV (Table I). For dimethylacetylene one also finds a near perfect matching of the experimental $IP_{\pi}^{\nu} = 9.61$ eV with the value 9.62 eV, calculated analogously by inserting the monomethylacetylene parameters $\epsilon_{\text{C}\equiv\text{C}}(\pi) = 11.40$ eV and $\epsilon_{\text{CH}_3}(\pi) = 14.25$ eV. The second and third π -orbitals are then expected at about 14.25 eV (nonbonding) and at about 16.18 eV (bonding), the other value obtained from the above calculation.

The results of the parameterized hyperconjugation model are displayed in Figure 3, together with the vertical ionization energies of those pe bands, which will be assigned to the $e(\pi)$ orbitals.

The assignment of the less well-resolved part of the disilyl- and dimethylacetylene pe spectra starts advantageously from the three $e(\pi)$ orbitals (Figure 3). As the lowest π ionization potential is well reproduced by the parameterization, the two higher ones are also considered trustworthy. Modified CNDO/2 calculations yield the orbital sequence $3e < 3a_1 < 2e < 1e < 2a_2 < 2a_1 < 1a_2 < 1a_1$ (cf. Figure 4).

Although intensity ratios are generally⁶ considered a rather poor guideline in assigning pe spectra, the linearly corrected values for the first three bands of disilylacetylene approaching 2:5:2 would fit to the CNDO/2 sequence as follows (cf. Table I): pe band 1, $3e$; shoulder 2, $3a_1$; multicomponent band 3, $2e$ plus one of the Jahn-Teller split $1e$; and band 4, the other $1e$ and $2a_2$. Obviously, there is no direct confirmation, but the vibrational fine structure on the low energy flank of the second pe band (Figure 1, shoulder 2) with $\bar{\nu} \sim 930$ cm^{-1} , probably due to the a_1 silyl deformation,⁸ would not be expected for a degenerate state with numerous overlapping fine structures. The 0.8 eV Jahn-Teller split observed for monosilylacetylene (Table I) lends some credit to the assumed $1e$ splitting. The remaining two bands at 18.2 and 19.0 eV are fitted into the above CNDO/2 sequence as $2a_1$ and $1a_2$.

The pe spectrum of dimethylacetylene (Figure 1) shows altogether only three separated bands. According to modified CNDO/2 calculations, the components 2-5 (Table I) of the second pe mountain




Figure 4. Diagrams showing main bond contributions to the occupied orbitals of disilylacetylene (according to modified CNDO/2 calculations for D_{3d} conformation).

stretching from 13.5 to 16.5 eV are in the same sequence, $3a_1 < 2e < 1e < 2a_2$, as in the disilyl derivative. Vibrational fine structures occur only in the first band ($3e(\pi)$: ground state, 2313 cm^{-1} ; ¹⁵ ion, 2050 cm^{-1}) and in the third band at 20.63 eV ($2a_1(?)$: ground state, $\nu_3^{\text{CH}_3}(a_1) = 1379 \text{ cm}^{-1}$; ¹⁵ ion, 1090 cm^{-1}). Thus the hyperconjugation model proposed for dimethylacetylene is in agreement with the correct prediction of the π band.

Calculations with or without π Interaction and Silicon 3d Basis Orbitals

In a simple LCBO MO hyperconjugation model, which uses only occupied orbitals as a basis, the interaction parameters $\beta_{\pi}^{\text{LCBO}}$ absorb the overall interaction, which cannot be subdivided any further on the basis of the available experimental data. To gain some insight into the interplay of those formal effects, *i.e.*, inductive and/or hyperconjugative destabilization *vs.* stabilizing back-bonding, LCAO MO calculations with and without π interactions and silicon 3d orbitals in the basis set have been performed.

For the calculations a modified CNDO/2 version (see Experimental Section) has been used, which in contrast to conventional CNDO/2 reproduces the spectroscopically deduced orbital sequences in both alkyl- and silylacetylenes, mainly due to the reduced π overlap the modified CNDO/2 readjusts, *e.g.*, the order $\pi(3e) < \sigma(3a_1)$ as observed experimentally. The calculations without any π interactions were achieved

(15) Cf. G. Herzberg, "Molecular Spectra and Molecular Structure," Vol. 2, Van Nostrand, New York, N. Y., 1945, p 356.




Figure 5. Modified CNDO/2 eigenvalues for the π orbitals of methylacetylene with or without π interaction.

by setting the off-diagonal elements between $\pi_{\text{C}\equiv\text{C}}$ and π_{XH_3} in the Hartree-Fock matrix equal to zero before diagonalization. The resulting "group orbital" energy $\pi_{\text{C}=\text{C}}(\text{H}_3\text{XC}\equiv\text{C}-)$, when compared with the calculated π orbital energy of acetylene, reflects the inductive effect of the XH_3 substituent. The procedure is illustrated in Figure 5 for methylacetylene. Within the above approximation the hyperconjugative destabilization of the acetylene π -orbital by a methyl group is smaller than the inductive one. In dimethylacetylene, hyperconjugation contributes 1 eV to the orbital shift and the inductive effect about 1.7 eV, the methyl group perturbations being by far not additive.

For silylacetylene two sets of calculations were carried out, without and with inclusion of silicon 3d orbitals (Figure 6). Comparison within each of the two sets reveals: Only the calculations with the reduced basis ($E_{\text{total}} = -327.2 \text{ eV}$) yield inductive as well as hyperconjugative π destabilization by the silyl group, both being smaller than the methyl group effects (Figure 5). Inclusion of the silicon 3d orbitals in the basis lowers the total energy as expected ($E_{\text{total}} = -334.7 \text{ eV}$) but leads to a "negative" hyperconjugative split $3.20 \rightarrow 3.05 \text{ eV}$ (Figure 6), shifting the $\pi_{\text{C}=\text{C}}$ orbital to higher values $|\epsilon_1|$ than in acetylene itself. This is contrary to the *pe* data (Table I), which clearly demonstrate, that the acetylene π ionization potential is lowered on silyl substitution. The reason for the exaggerate $p_{\pi}d_{\pi}$ back-bonding is traced back to unrealistic overlap relations

$$\begin{array}{cc} \text{H}_3\text{C} \text{---} \text{C}(\equiv\text{CH}) & \text{H}_3\text{Si} \text{---} \text{C}(\equiv\text{CH}) \\ 1.46 \text{ \AA} & 1.83 \text{ \AA} \\ S(p_{\pi}p_{\pi}) = 0.13 & S(p_{\pi}p_{\pi}) = 0.12 \\ & S(p_{\pi}d_{\pi}) = 0.18 \end{array}$$

The much too high overlap $S(p_{\pi}d_{\pi})$ leads to a Hartree-Fock off-diagonal element of 2.6 eV compared to a $p_{\pi}p_{\pi}$ interaction of only 1.7 eV. As expected from the overlap disproportion, the CNDO/2 calculations with inclusion of $3d_{\text{Si}}$ orbitals also fail for disilylacetylene, while those with the reduced $3s_{\text{Si}}/3p_{\text{Si}}$ basis set allow comparison with other acetylenes as well as with the *pe* data. Analogous comments were first published by

Pitt,¹⁶ who recently, in an extensive examination⁴ of available experimental data, concluded that numerous ground-state properties of group IV compounds are reasonably reproduced by CNDO/2 calculations with sp basis set. This statement is further confirmed by the calculated charge distributions and dipole moments of methyl- and silylacetylenes. Convincingly, trends and magnitudes of the experimental gas-phase dipole moments¹⁷ are only reflected in modified CNDO/2 calculations without silicon 3d orbitals.

	without	with 3d _{Si}
$\text{H}_3\text{C-C}\equiv\text{C-H}$	3(+0.038)	3(-0.074)
$\text{H}_3\text{Si-C}\equiv\text{C-H}$	3(-0.074)	3(-0.100)
μ^{calcd}	1.013 D	0.135 D
μ^{exptl}	0.781 D ¹⁷	0.316 D ¹⁷

Concluding Remarks

The preceding discussions of methyl- and silylacetylenes and especially their pe spectra have demonstrated: (i) how simple LCBO MO hyperconjugation models can be parameterized using pe data and how well they predict ionization potentials of related compounds, (ii) why LCAO MO calculations using modified CNDO/2 exaggerate p_πd_π back-bonding, and (iii) that a basis set without silicon d orbitals better reproduces the dipole moment of silylacetylene and only allows comparison of pe ionization potentials and CNDO/2 eigenvalues within the series RC≡CR (R = H, CH₃, SiH₃).

The main concern for the chemist might be not so much whether 3d_{Si} participation is real and provable or not but rather which model will be most useful and easily applicable to rationalize experimental data. Our example shows, that the different lowering of the acetylene π ionization potentials by methyl and silyl groups can be explained by both LCAO MO and LCBO MO models without explicit introduction of p_πd_π back-bonding. No decision can be made, however, on how much of it might be absorbed in the parameterization—especially within an LCBO MO hyperconjugation approach, which only considers occupied orbitals.

Cradock, Ebsworth, and Whiteford,¹⁰ who published first the pe spectra of most of the silyl and germyl derivatives, E(XH₃)_n, of group V to VII elements E, based their assignments mainly on orbital symmetry considerations. Cautiously they pointed out, that the following observations might be in accord with d orbital participation:¹⁰ (i) the first ionization potentials of compounds E(XH₃)_n always reach their maximum value in the silyl derivative $IP_{\text{CH}_3} < IP_{\text{SiH}_3} > IP_{\text{GeH}_3}$, (ii) spin-orbit coupling of heavy atoms is reduced in the silyl relative to the methyl derivative, and (iii) pe bands assigned to electron pair ionizations are usually broadened in the spectra of silyl compounds. These findings could be rationalized within an LCBO MO model as follows. (i) The maximum ionization potential $IP_{\text{CH}_3} <$

(16) C. G. Pitt, *Chem. Commun.*, 816 (1971); *J. Organometal. Chem.*, 23, C 35 (1970).

(17) R. D. Nelson, D. R. Lide, and A. A. Maryott, "Selected Values of Electric Dipole Moments," National Bureau of Standards NSRDS-NBS, 1967, p 10.





Figure 6. Modified CNDO/2 eigenvalues for the π-orbitals of silylacetylene with and without π overlap and silicon 3d orbitals.

$IP_{\text{SiH}_3} > IP_{\text{GeH}_3}$ needs two opposing effects for its explanation, e.g., different inductive electron donation and counteracting p_πd_π back-bonding in the sequence CH₃-SiH₃-GeH₃. Within the hyperconjugation model, analogous second-order perturbation arguments can be employed; the change in orbital energies is proportional to the square of the interaction term β² and thus depends on overlap respectively on bond lengths and inversely proportional to the energy difference between the coulomb parameters Δα and thus depends on the relative position of the basis orbitals. Therefore, depending on both the parameters, α and β, a minimum eigenvalue ε_i, i.e., via Koopmans' theorem -ε_i ≡ IP_n, a maximum first-ionization potential can be reproduced. (ii) To incorporate second-order spin-orbit coupling qualitatively into an LCBO MO model,¹⁸ symmetry double groups¹⁹ prove to be useful. For a C_{3v} molecule as silyl iodide,¹⁰ H₃SiI, on inclusion of spin functions in the C_{3v}' double group, only two irreducible representations result, e_{3/2} and e_{1/2}. Symmetry allowed mixing yields



Obviously, the splitting between the adjacent e_{1/2} orbitals must exceed that between the more distant e_{3/2} ones. Thus for silyl iodide a smaller gap between the first two pe bands is expected than for methyl iodide, judging along from second-order spin-orbit coupling. (iii) Also the reasoning for the observed broadening⁶ of the first pe bands of silyl compounds¹⁰ does not necessarily have to rest on p_πd_π delocalization. Within an LCBO MO model it is as easily explained by the stronger mixing of two symmetry equivalent orbitals as their energy distance Δα shortens.

Repeatedly it has been pointed out that there is no chance to subdivide LCBO MO parameters unequivocally any further on the basis of experimental data.

(18) K. Wittel, H. Bock, and R. Manne, *Tetrahedron*, in press; K. Wittel, *Chem. Phys. Lett.*, 15, 555 (1972).

(19) Cf. for instance G. Herzberg, "Molecular Spectra and Molecular Structure," Vol. III, Van Nostrand, New York, N. Y., 1966, p 563.

Therefore the hyperconjugation model, well suited to help in assigning the π spectra of alkyl- and silylacetylenes, does neither prove nor disprove $p_x d_x$ back-bonding in silicon compounds—just as most of the numerous arguments produced favor one or the other.

Experimental Section

Synthesis. Monosilylacetylene²⁰ was prepared from acetylene magnesium bromide and iodasilane in dry, degassed tetrahydrofuran. The reaction product was pumped off and purified by fractionate condensation. The measurement sample was finally obtained by high vacuum line distillation between two baths at -112 and -132° and its purity controlled by ir²¹ and nmr²⁰ as well as molecular weight determination (found 56.8, calcd 56.1).

Disilylacetylene²² can be obtained analogously from acetylene di-Grignard solution in dibutyl ether and bromosilane. The measurement sample was purified from dibutyl ether by fractionate

condensation at -78 and at -96° from excess bromosilane. Neither by ir and nmr nor by molecular weight determination (found 87.0, calcd 86.2) could any impurities in the measurement sample be detected.

Spectra. The π spectrum of methylacetylene was taken from the literature;⁷ the one of dimethylacetylene has been kindly given to us by Elbel²³ from our group. The π spectra of the silylacetylenes were recorded with a Perkin-Elmer PS 16 and calibrated with argon. The samples have been introduced into the spectrometer carefully avoiding exposition to air.

Calculations. The calculations have been carried out at the UNIVAC 1108 of the computer center of the University of Frankfurt. The modified CNDO/2 version used is described elsewhere.²⁴ In addition the β_A^0 values for the third-period elements have been reduced proportionally to β_{carbon}^0 .²⁴ The π interaction was switched off by setting the corresponding $A(i,j)$ in subroutine SCFCLO equal to zero. Silicon d orbitals can be excluded by either reducing the basis set or by setting all d overlap integrals in subroutine INTEGRL equal to zero. The latter procedure allows alternatively a reduction of the back-bonding by multiplying the d overlap integrals with a factor <1 .

(20) E. A. V. Ebsworth and S. G. Frankiss, *J. Chem. Soc.*, 661 (1963).

(21) E. A. V. Ebsworth, S. G. Frankiss, and W. J. Jones, *J. Mol. Spectrosc.*, **13**, 9 (1964); R. B. Reeves, R. E. Wilde, and D. W. Robinson, *J. Chem. Phys.*, **40**, 125 (1964).

(22) R. C. Lord, D. W. Mayo, H. E. Opitz, and J. S. Peake, *Spectrochim. Acta*, **12**, 147 (1958).

(23) Part of the thesis of S. Elbel, unpublished results.

(24) J. Kroner, D. Proch, W. Fuss, and H. Bock, *Tetrahedron*, **28**, 1585 (1972).